

QUANTUM SUPERMEMBRANES AND ADS/CFT DUALITY

A. A. Tseytlin^{a,b,c*}^a *Blackett Laboratory, Imperial College
SW7 2AZ, London, UK*^b *Institute for Theoretical and Mathematical Physics, Moscow State University
119991, Moscow, Russia*^c *Lebedev Physical Institute, Russian Academy of Sciences
119991, Moscow, Russia*Received September 5, 2023,
revised version September 5, 2023
Accepted for publication September 7, 2023

Extended Abstract. The full text of this paper is published in the English version of JETP.

DOI: 10.31857/S0044451023120118

EDN: MXMLBW

Dedicated to the memory of Valery Rubakov

Testing AdS/CFT duality [1, 2] is an important direction of research leading to novel non-trivial results on both gauge theory and dual string theory sides. Most of the work so far was done for duality of 4d super Yang-Mills theory and type IIB 10d string theory in $AdS_5 \times S^5$ background. The aim of this paper is to review and extend some recent work [3, 4] on testing AdS/CFT correspondence between supersymmetric $U(N)_k \times U(N)_{-k}$ Chern-Simons-matter 3d gauge theory and M-theory in $AdS_4 \times S^7/\mathbf{Z}_k$ background [5, 6] (ABJM theory).

Membrane theory with the bosonic action dating back to Dirac [7] remains an enigma. While the existence of a consistent quantum theory of bosonic membranes may be in doubt, it may happen to be well defined for the 11d supermembrane or M2 brane theory [8]. The large amount of supersymmetry and possibly some unknown hidden symmetries may lead to its UV finiteness despite formal power-counting nonrenormalizability. This may be true, in particular, for the supermembrane in the maximally supersymmetric $AdS_4 \times S^7$ or $AdS_7 \times S^4$ backgrounds [9, 10].

While the M2 brane action is highly non-linear, expanded near a classical solution with non-degenerate

induced 3d metric it can be quantized in a static gauge. Then the leading 1-loop result for its partition function is UV finite and thus unambiguous [3, 4, 11, 12].

The M2 brane action in 11d background is related to the type IIA string in the corresponding 10d background by a double dimensional reduction [13]. Considering M2 brane world volume of topology $\Sigma^2 \times S^1$ and expanding 3d fields in Fourier modes in S^1 coordinate one gets an effective 2d string action on Σ^2 coupled to an infinite tower of massive 2d fields. Choosing a static gauge in the M2 brane action one gets a static gauge Nambu-Goto action for the massless transverse string modes coupled to a tower of the massive “Kaluza-Klein” 2d modes. This “effective string” 2d action is essentially equivalent to the original M2 brane action and thus may inherit some of its hidden symmetries.

The work described below provides a remarkable evidence that direct semiclassical quantization of the M2 brane in $AdS_4 \times S^7/\mathbf{Z}_k$ background reproduces the results of large N localization computations [14–16] of the 1/2-BPS Wilson loop and the instanton contributions to free energy in the ABJM gauge theory.

Our first example is the 1/2-BPS Wilson loop. For fixed k , the large N expansion of the Wilson loop operator in the ABJM theory corresponds to the expansion in the large effective M2 brane tension $R^3 T_2 \sim \sqrt{Nk}$, where R is the curvature radius of $AdS_4 \times S^7/\mathbf{Z}_k$ and $T_2 = 1/((2\pi)^2 \ell_P^3)$. The analytic expression for the expectation value of the 1/2-BPS circular Wilson loop in the ABJM theory was derived

* E-mail: atseytlin@gmail.com

using supersymmetric localization in [15]. In order to compare to the semiclassical expansion in the M2 brane world-volume theory, one is to expand this expression at large N with fixed k , which gives

$$\langle W_{\frac{1}{2}} \rangle = \frac{1}{2 \sin(\frac{2\pi}{k})} e^{\pi \sqrt{\frac{2N}{k}}} \times \left[1 - \frac{\pi(k^2 + 32)}{24\sqrt{2}k^{3/2}} \frac{1}{\sqrt{N}} + O\left(\frac{1}{N}\right) \right]. \quad (1)$$

The Wilson loop has a dual description in terms of an M2 brane wrapped on $\text{AdS}_2 \times S^1$ [17] in the M-theory background $\text{AdS}_4 \times S^7/\mathbf{Z}_k$. One finds [3] that the exponential factor in (1) is reproduced by the classical action of the M2 brane with $\text{AdS}_2 \times S^1$ world-volume, while the k -dependent prefactor $(2 \sin(2\pi k))^{-1}$ is matched precisely by the 1-loop correction coming from the functional determinants of the quantum fluctuations around this M2 brane solution.

The non-perturbative part of the ABJM theory free energy found from localization is [16]

$$F^{\text{np}}(N, k) = F_1^{\text{inst}}(N, k) \times \left[1 + \frac{\pi}{\sqrt{2k}} \frac{k^2 - 40}{12k} \frac{1}{\sqrt{N}} + \dots \right], \quad (2)$$

$$F_1^{\text{inst}}(N, k) = -\frac{1}{\sin^2(\frac{2\pi}{k})} e^{-2\pi \sqrt{\frac{2N}{k}}}.$$

Here F_1^{inst} is the leading large N term in the 1-instanton contribution. In the type IIA string theory regime (i.e., for large N and k with $N/k = \text{fixed}$) it may be interpreted as the contribution of the string world-sheet instanton (wrapping CP^1 in CP^3 [18]). In the M-theory regime (i.e. for large N with fixed k), the world-sheet instantons correspond to the M2 brane instantons wrapping the 11d circle and a CP^1 in CP^3 , i.e., $S^3/\mathbb{Z}_k \subset S^7/\mathbf{Z}_k$. Computing the corresponding classical M2 brane action and 1-loop fluctuation determinants [4] one matches precisely the exponential and its prefactor in (2).

As another example of the quantum M2 brane computation in $\text{AdS}_4 \times S^7/\mathbf{Z}_k$ one may consider the 1-loop correction to the partition function expanded near the classical M2 brane solution generalizing infinitely long rotating folded string [19, 20] in AdS_4 . It determines leading large λ corrections at each order in $1/N$ to the cusp anomaly function in ABJM theory [21]. The predicted generalization of the planar cusp anomaly in

ABJM theory [22–24] to include leading at large terms at each order in $1/N$ reads

$$\hat{f}(k) = c_0 + f(k) = -\frac{5}{2\pi} \log 2 + \frac{2\pi}{3k^2} + \frac{2\pi^3}{45k^4} + \dots = -\frac{5}{2\pi} \log 2 + \frac{2\pi\lambda^2}{3N^2} + \frac{2\pi^3\lambda^4}{45N^4} + \dots$$

From the point of view of the 't Hooft limit of large N and large $k = N/\lambda$ these corrections are non-planar corrections to the cusp anomaly. This result is a non-trivial strong-coupling prediction about non-planar correction to the anomalous dimension of a high-spin operator in the ABJM theory. Being non-planar it is unlikely to be captured by integrability methods. Also, as the cusp anomaly corresponds to a non-supersymmetric observable (anomalous dimension of a non-BPS operator) it is unlikely to be captured by a localization computation.

The full text of this paper is published in the English version of JETP.

REFERENCES

1. J. M. Maldacena, *The Large N Limit of Superconformal Field Theories and Supergravity*, Adv. Theor. Math. Phys. **2**, 231 (1998), <https://doi.org/10.4310/ATMP.1998.v2.n2.a1>, <https://arxiv.org/abs/hep-th/9711200>.
2. O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri, and Y. Oz, *Large N Field Theories, String Theory and Gravity*, Phys. Rept. **323**, 183 (2000), [https://doi.org/10.1016/S0370-1573\(99\)00083-6](https://doi.org/10.1016/S0370-1573(99)00083-6), <https://arxiv.org/abs/hep-th/9905111>.
3. S. Giombi and A. A. Tseytlin, *Wilson Loops at Large N and the Quantum M2-Brane*, Phys. Rev. Lett. **130**, 201601 (2023), <https://doi.org/10.1103/PhysRevLett.130.201601>, <https://arxiv.org/abs/2303.15207>.
4. M. Beccaria, S. Giombi, and A. A. Tseytlin, *Instanton Contributions to the ABJM Free Energy from Quantum M2 Branes*, <https://arxiv.org/abs/2307.14112>.
5. O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, *$N = 6$ Superconformal Chern–Simons–Matter Theories, M2-branes and their Gravity Duals*, JHEP **10**, 091 (2008), doi.org/10.1088/1126-6708/2008/10/091, <https://arxiv.org/abs/0806.1218>.
6. J. Bagger, N. Lambert, S. Mukhi, and C. Papageorgakis, *Multiple Membranes in M-theory*, Phys. Rept. **527**, 1 (2013), <https://doi.org/10.1016/j.physrep.2013.01.006>, <https://arxiv.org/abs/1203.3546>.

7. P. A. M. Dirac, *An Extensible Model of the Electron*, Proc. Roy. Soc. Lond. A **268**, 57 (1962), <https://doi.org/10.1098/rspa.1962.0124>.
8. E. Bergshoeff, E. Sezgin, and P. K. Townsend, *Supermembranes and Eleven-Dimensional Supergravity*, Phys. Lett. B **189**, 75 (1987), [https://doi.org/10.1016/0370-2693\(87\)91272-X](https://doi.org/10.1016/0370-2693(87)91272-X).
9. E. Bergshoeff, M. J. Duff, C. N. Pope, and E. Sezgin, *Compactifications of the Eleven-Dimensional Supermembrane*, Phys. Lett. B **224**, 71 (1989), [https://doi.org/10.1016/0370-2693\(89\)91053-8](https://doi.org/10.1016/0370-2693(89)91053-8).
10. B. de Wit, K. Peeters, J. Plefka, and A. Sevrin, *The M Theory Two-Brane in $AdS_4 \times S^7$ and $AdS_7 \times S^4$* , Phys. Lett. B **443**, 153 (1998), [https://doi.org/10.1016/S0370-2693\(98\)01340-9](https://doi.org/10.1016/S0370-2693(98)01340-9), ArXiv hep-th/9808052.
11. M. J. Duff, T. Inami, C. N. Pope, E. Sezgin, and K. S. Stelle, *Semiclassical Quantization of the Supermembrane*, Nucl. Phys. B **297**, 515 (1988), [https://doi.org/10.1016/0550-3213\(88\)90316-1](https://doi.org/10.1016/0550-3213(88)90316-1).
12. N. Drukker, S. Giombi, A. A. Tseytlin, and X. Zhou, *Defect CFT in the 6d (2,0) Theory from M2 Brane Dynamics in $AdS_7 \times S^4$* , JHEP **07**, 101 (2020), [https://doi.org/10.1007/JHEP07\(2020\)101](https://doi.org/10.1007/JHEP07(2020)101), <https://arxiv.org/abs/2004.04562>.
13. M. J. Duff, P. S. Howe, T. Inami, and K. S. Stelle, *Superstrings in $D=10$ from Supermembranes in $D=11$* , Phys. Lett. B **191**, 70 (1987), [https://doi.org/10.1016/0370-2693\(87\)91323-2](https://doi.org/10.1016/0370-2693(87)91323-2).
14. N. Drukker, M. Marino, and P. Putrov, *From Weak to Strong Coupling in ABJM Theory*, Commun. Math. Phys. **306**, 511 (2011), <https://doi.org/10.1007/s00220-011-1253-6>, <https://arxiv.org/abs/1007.3837>.
15. A. Klemm, M. Marino, M. Schiereck, and M. Soroush, *Aharony–Bergman–Jafferis–Maldacena Wilson Loops in the Fermi Gas Approach*, Z. Naturforsch. A **68**, 178 (2013), <https://doi.org/10.5560/ZNA.2012-0118>, <https://arxiv.org/abs/1207.0611>.
16. Y. Hatsuda, S. Moriyama, and K. Okuyama, *Instanton Effects in ABJM Theory from Fermi Gas Approach*, JHEP **01**, 158 (2013), [https://doi.org/10.1007/JHEP01\(2013\)158](https://doi.org/10.1007/JHEP01(2013)158), <https://arxiv.org/abs/1211.1251>.
17. M. Sakaguchi, H. Shin, and K. Yoshida, *Semiclassical Analysis of M2-brane in $AdS_4 \times S^7/Z_k$* , JHEP **12**, 012 (2010), [https://doi.org/10.1007/JHEP12\(2010\)012](https://doi.org/10.1007/JHEP12(2010)012), <https://arxiv.org/abs/1007.3354>.
18. A. Cagnazzo, D. Sorokin, and L. Wulff, *String Instanton in $AdS(4) \times CP^3$* , JHEP **05**, 009 (2010), [https://doi.org/10.1007/JHEP05\(2010\)009](https://doi.org/10.1007/JHEP05(2010)009), <https://arxiv.org/abs/0911.5228>.
19. S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *A Semiclassical Limit of the Gauge/String Correspondence*, Nucl. Phys. B **636**, 99 (2002), [https://doi.org/10.1016/S0550-3213\(02\)00373-5](https://doi.org/10.1016/S0550-3213(02)00373-5), <https://arxiv.org/abs/hep-th/0204051>.
20. S. Frolov and A. A. Tseytlin, *Semiclassical Quantization of Rotating Superstring in $AdS(5) \times S^5$* , JHEP **06**, 007 (2002), <https://doi.org/10.1088/1126-6708/2002/06/007>, <https://arxiv.org/abs/hep-th/0204226>.
21. S. Giombi and A. Tseytlin, Unpublished.
22. T. McLoughlin and R. Roiban, *Spinning Strings at One-Loop in $AdS(4) \times P^3$* , JHEP **12**, 101 (2008), <https://doi.org/10.1088/1126-6708/2008/12/101>, <https://arxiv.org/abs/0807.3965>.
23. L. F. Alday, G. Arutyunov, and D. Bykov, *Semiclassical Quantization of Spinning Strings in $AdS(4) \times CP^3$* , JHEP **11**, 089 (2008), <https://doi.org/10.1088/1126-6708/2008/11/089>, <https://arxiv.org/abs/0807.4400>.
24. T. McLoughlin, R. Roiban, and A. A. Tseytlin, *Quantum Spinning Strings in $AdS(4) \times CP^3$: Testing the Bethe Ansatz Proposal*, JHEP **11**, 069 (2008), <https://doi.org/10.1088/1126-6708/2008/11/069>, <https://arxiv.org/abs/0809.4038>.